

# **A Study of Equine Ulcers and Dietary Supplements**

Ulcers in horses are ubiquitous, but a unique formulation of natural ingredients can reduce their size and number.

Last Major Revision:

November 19, 2015

Updated July 26, 2017

# Introduction

Franklin L. Pellegrini, DVM and Scott C. Anderson

Studies over the past decade have documented an unsettling fact: over 90% of race horses and 60% of show horses have gastric ulcers.<sup>1</sup> Worse yet, the number one killer of horses is colic,<sup>2</sup> which is an expensive and unnecessary tragedy.

This unfortunate state of affairs, known as equine gastric ulcer syndrome (EGUS), is due to the heavy training and unnatural feeding regimens employed to maximize performance. Horses in the wild graze in herds all day and only sporadically exert themselves, but stabled race and show horses are kept in confinement and fed high carbohydrate feed two or three times a day in order to increase their performance. Carbohydrate loads in excess of what the small intestine can absorb continue to the hindgut, where a portion of them are fermented by bacteria into lactic acid, which can lead to hindgut acidosis.<sup>3</sup>

Unlike humans who secrete digestive juices on demand, horses secrete stomach acids on a fairly continuous basis. As they eat, they swallow bicarbonate-containing saliva at the rate of five gallons a day, helping to neutralize these stomach acids. When they eat only two or three times a day, there is far less saliva to buffer the gastric juices, which include hydrochloric acid, bile and pepsin, and can reach a highly acidic pH of 1.5. Research done at the University of Florida has shown that heavy training may then move these acids around to less protected parts of the stomach.<sup>4</sup>

This establishes an unnatural stomach environment that may ultimately lead to ulceration. Horse ulcers are most prevalent in the upper, squamous area of the stomach and the *margo plicatus*, but they are often found in the glandular lower section as well.<sup>5</sup> These ulcers in turn lead to loss of blood, irritability and poor absorption of nutrients. Paradoxically, this may negatively impact the performance of exactly those horses which are expected to work at peak efficiency. Ulcers may be responsible for much of the subclinical anemia, listlessness, weight loss and general poor health noted by many performance horse veterinarians for years. In fact, the number one complaint about horses with ulcers is that their performance is declining.<sup>6</sup> With these animals, environment has a much larger impact on their ability than genetic factors.<sup>7</sup>

Gastric ulcers in horses can be inspected by endoscopy, but colonic ulcers are much harder to observe. Colonoscopies are impractical due to the difficulty of evacuating the equine colon without endangering the health of the horse. Lacking evidence to the contrary, it has generally been felt that equine colonic ulcers are rare.

This paper reports on two recent studies that cast a worrying light on equine ulcers, demonstrating that in addition to gastric ulcers, horses are also plagued by colonic ulcers, afflicting over 60% of performance horses.

Additionally, one of these studies indicates that the most commonly prescribed drugs for treating gastric ulcers has little or no effect on colonic ulcers. Finally, this paper discusses a novel feed supplement that has been shown to reduce both gastric and colonic ulcers in performance horses.

## References

---

- <sup>1</sup> McClure SR, Glickman LT, Glickman NW. *Prevalence of gastric ulcers in show horses*. J Am Vet Med Assoc 1999;215:1130-1133.
- <sup>2</sup> Traub-Dargatz JL, Koprak CA, Seitzinger AH, Garber LP, Forde K, White NA. *Estimate of the national incidence of and operation-level risk factors for colic among horses in the United States, spring 1998 to spring 1999*. J Am Vet Med Assoc. 2001 Jul 1;219(1):67-71.
- <sup>3</sup> Richards, N, Gn Hinch, and Jb Rowe. “The Effect of Current Grain Feeding Practices on Hindgut Starch Fermentation and Acidosis in the Australian Racing Thoroughbred.” Australian Veterinary Journal 84, no. 11 (2006): 402–407. doi:10.1111/j.1751-0813.2006.00059.x.
- <sup>4</sup> Lorenzo-Figueras M, Merritt AM. *Effects of exercise on gastric volume and pH in the proximal portion of the stomach of horses*. Am J Vet Res. 2002;63:1481-1487]
- <sup>5</sup> Mair, Tim, Tom Divers, and Norm Ducharme. *Manual of Equine Gastroenterology*. London: WB Saunders, 2002.
- <sup>6</sup> Mitchell RD. *Prevalence of Gastric Ulcers in Hunter/Jumper and Dressage Horses Evaluated for Poor Performance*. Assoc. Equine Sports Med., September 2001.
- <sup>7</sup> Kronfeld DS. *Speed Limit*. Dipl. ACVN, Dipl. ACVIM. March 2003 Article # 4212.

## Functional Feeding and Equine GI Health

By Peter M. Bedding, PhD and Scott C. Anderson

Feeds are not medicines but good feed management is probably the best long-term solution to many digestive problems, including ulcers. Ironically, feed is also the main culprit in many of these disorders. This is because trainers often put their horses on hard feed in order to guarantee high energy concentrations for maximum performance. But in addition to giving them extra energy, hard feeds also allow large quantities of sugars to be released in the mouth.

Horses have evolved to eat large quantities of grass, and in the process they produce gallons of saliva daily. But when it is allowed to chew only small amounts of hard feed, a performance horse doesn't generate enough saliva to buffer the high levels of sugars and acids that are produced.

As well as decaying their teeth, the hard feed adds to the acidity of the stomach, which in due course challenges the stomach wall. Together with stress and exercise, this can lead to stomach ulcers.

A natural diet supplies substances that support the integrity of the GI tract. However, in emphasizing performance, many manufactured feedstuffs end up being deficient in naturally occurring nutrients that are needed to maintain the health of a horse.

Vitamins and minerals are typically added in an attempt to correct these shortcomings. When various nutrients are compounded to address a specific function – such as immune response or tendon healing – it is called functional feeding, and the formulation is called a nutraceutical. But before any functional feeds can have an effect, the gut itself must be in good shape. Therefore, a nutraceutical that addresses the proper functioning of the gut provides a necessary foundation for all other functional feeds.

Such a gut-oriented functional feed would include specific prebiotics, amino acids, nucleotides, polar lipids and antioxidants. With a nutraceutical like this, the health of the GI tract can be actively managed, protecting the gut wall from attack by acids, pathogens and toxins and promoting the regeneration of intestinal tissue cells.

Pasturing a horse is a sure-fire way to improve its digestion, but it doesn't provide the energy density needed for a performance horse. In the absence of the ideal natural feeding environment, a functional feed is essential.

In today's high performance horse world, ulcers are ubiquitous and colic is the number one killer. But with the proper functional feeding program, the equine GI tract can quickly recover from insults even during strenuous training, allowing the horse to operate at the absolute peak of its abilities.

## A Large-Scale Necroscopic Study

Franklin L. Pellegrini, DVM and Scott C. Anderson


In 2004, one of the authors (Pellegrini) conducted a large-scale necropsy on 180 performance horses in a Texas abattoir. It was found that 87% of these horses had gastric ulcers and 63% had colonic ulcers, resulting in an overall ulceration rate of 97%.<sup>1</sup> These results were then correlated to a guaiac-based Fecal Occult Blood Test (gFOBT).

After being euthanized, necropsies were performed on the horses. Gastric ulcers were observed and graded on a standard scale of 0-4, as described below:

- **Grade 0:** Normal, unulcerated tissue. The epithelium is intact and there is no thickening or abnormal coloring.
- **Grade 1:** The mucus lining is intact, but there are areas of thickened, discolored tissue.
- **Grade 2:** Small, single or multiple ulcers present.
- **Grade 3:** Large, single or multiple ulcers.
- **Grade 4:** Extensive, deep ulcers.

Colonic ulcers were also observed and – lacking an accepted standard of measurement – were marked as either extant or not.

The results may be summarized by the following graph:



**Incidence of gastric and colonic ulcers. Note the large overlap representing horses with both gastric and colonic ulcers. Only 3% of horses have no ulcers.**

Prior to the dissection of the horse, a fecal sample was collected. This manure was tested with guaiac and correlated to the gross examination of intestinal tissue.

Overall, the guaiac test proved to be highly specific and significant (100% for both, with P=0) for the existence of an ulcer, due to the unexpected lack of false positives in this group. The existence of false negatives, however, lowered the overall accuracy of the test to 65%, roughly comparable to human outcomes with such a test.

Guaiaac works by binding hemoglobin and turning blue in the presence of hydrogen peroxide. The false negatives are likely due to the fact that the guaiac reaction is not very sensitive, requiring 10mg of hemoglobin per gram of stool to produce a positive result. Since many ulcers bleed only lightly or sporadically, these may be missed by the gFOBT.

Importantly, this study demonstrated that when there is a positive gFOBT in the absence of gastric ulcers, then colonic ulcers are always present. Thus, endoscopy in combination with a gFOBT can provide a significant test for colonic ulcers.

---

<sup>i</sup> Pellegrini, FL. *Results of a large-scale necroscopic study of equine colonic ulcers*. J Equine Vet Sci. 2005;25(3):113-17.

# A Comparative Study

By Franklin L. Pellegrini, DVM and Scott C. Anderson

In order to evaluate the effectiveness of a novel feed supplement containing oat- and yeast-based nutraceuticals, an 80-horse study was conducted at an Ohio Standardbred racehorse stable. For the test, 20 horses were assigned randomly to each of four groups, including a control, a group using proton-pump inhibitors and two groups that were administered different dosages of the feed supplement.

## Proton-Pump Inhibitors

Proton-pump inhibitors (PPIs) are a class of drugs that block the production of stomach acid. These are typically prescribed for horses that are known by endoscopy to have ulcers, but are also prescribed for non-specific ailments on the theory that if improvement is noted, then an ulcer was likely to blame.


In a blind test of 20 Standardbred horses on a proton-pump inhibitor vs. 20 controls, it was found that acid inhibition was effective at reducing gastric ulcer severity (graded on a scale from 0-4) in the first month of treatment as determined by endoscopy.

This comparative study also looked at the effects of two different dosages of an oat- and yeast-based nutritional supplement (SUCCEED DCP) on the rate of ulceration. Using two other groups of 20 horses each, DCP showed a dosage-dependent improvement in gastric ulcers, reducing the average severity to less than 0.5 by the end of the study (error bars show standard error, adjusted for population size).


During the period of this study, a guaiac-based fecal blood test (gFOBT) of the manure was also conducted. The guaiac test can produce false positives when exposed to blood from eating red meat; fortunately this is not a problem for herbivores. Nevertheless, it has been suggested that the guaiac test might be skewed toward false positives in horses as well, perhaps due to chlorophyll or other peroxidase-mimicking molecule in their feed. However, in an independent (unpublished) Texas study we found both gastric and colonic ulceration correlated with a high significance and specificity to a positive guaiac stain.

Interpreted in this light, the gFOBT data can serve as a proxy for colonic and other non-gastric ulcers. The study demonstrated an overall dosage-dependent reduction in these non-gastric ulcers for horses using the nutritional supplement. Proton pump inhibitors, which are specifically designed for gastric ulceration, did not show a decrease over this time period.

### Gastric Ulcer Time Study



### gFOBT Time Study



These dosage-dependent results are a strong indicator of the efficacy of DCP for reducing fecal blood, a proxy for colonic ulcers in performance and race horses.

Further studies are planned, using a more sensitive ELISA (enzyme-linked immunosorbant assay) test, to better analyze the presence and extent of colonic ulcers.

## **Conclusions**

This paper has several findings:

- Not only do horses have gastric ulcers, they also have a high incidence of colonic ulcers.
- Proton-pump inhibitors are useful to quell gastric ulcers, but they have little effect on colonic ulcers.
- A nutritional feed supplement made with oat oil, oat flour, amino acids, nucleotides and yeast extracts including mannan oligosaccharides (SUCCEED DCP) has been shown to deliver a strong dosage-dependent reduction in both gastric and colonic ulcers.

## **Next Steps**

Further research is scheduled to investigate anecdotal reports of improved equine disposition, appearance, performance and reproductive success. In addition, a more sensitive ELISA test is being developed that will be used in a follow-up necroscopic study of equine ulceration.

# A Nutritive Supplement for Horses

Peter M.J. Bedding, PhD and Scott C. Anderson

In order to treat these gastric and colonic ulcers in horses, one of the authors (Bedding), devised a feed supplement to target each aspect of the problem. The hope was to create a digestive conditioning program that would normalize conditions in the GI tract while still allowing horses to be rigorously trained.

This feed supplement was subsequently tested on 80 horses in Ohio. The key ingredients of this patent-pending formula include:

## Polar Lipids

Lipids represent a large class of molecules that include fatty acids, phospholipids (lecithin), galactolipids and triglycerides. They play a key role in the structure and function of cellular membranes and are found in much of the plant material already in equine diets. As a consequence of their ubiquity, lecithins and lipids are considered to be a GRAS (generally regarded as safe) supplement.

Oat oil is rich in polar lipids, particularly galactolipids. These are rare in animals, but are the most common lipids in plants as well as the most abundant form of lipids on the planet. Animals generally lack the enzymes needed to synthesize these polyunsaturated polar lipids, and so they must acquire them in the diet. These dietary polar lipids are important in forming the tight junctions between the epithelial cells lining the gut. Cells connected in this fashion present a unified barrier against digestive juices, toxins and pathogens. When these junctions are disrupted, the defensive wall is breached, and injury in the form of an ulcer may result. Adding polar lipids to the diet has been shown to protect the intestinal mucosa and to strengthen the impermeability of the barrier.<sup>1 2 3</sup>

As well as galactolipids, oat oil is rich in trienols and tocopherols, powerful anti-oxidants that sweep free radicals out of the system before they can damage the surrounding tissue.

Polar lipids are versatile emulsifiers, stabilizing oil-water mixtures. They provide an ideal nutrient delivery vehicle, capable of ferrying both fat- and water-soluble molecules into the tissues. Fat-soluble nutrients include vitamins A, K, D and E. Polar lipids, especially galactolipids, have been shown to increase bio-availability of such nutrients by up to 500%.

After transporting their nutritive load, polar lipids are readily absorbed in the gut (after digestion by bile salts), where they supply extra energy to the horse.<sup>4</sup>

## Beta-glucan

Beta-glucan is a polysaccharide: a chain of glucose molecules that can branch in specific conformations, each one having unique properties. This branching is represented by a beta notation that describes how the chains are linked, such as  $\beta(1,3)$ . Different beta-glucans can be derived from yeast, barley and oats, and they can have profound effects on typical animal

systems. They have been known for years to reduce LDL cholesterol levels in the blood.<sup>5 6</sup>

More significantly, beta-glucans moderate the release of sugars from the digestive system, helping to prevent the sugar highs and lows that often afflict animals that are fed only two or three times a day on an energy-rich high-carbohydrate diet.<sup>7</sup> Studies have shown that beta-glucan is effective in reducing post-prandial glycemic peaks by up to 50%.<sup>8</sup>

Although more equine studies are needed, research has implicated bacteria in the formation of ulcers in many animals. While it has been difficult to culture bacteria from horse stomachs, colonic bacteria are generally known to include a wide variety of pathogenic species.

Beta-glucan is a potent stimulator of the immune system. It arouses macrophages – which have a specific beta-glucan receptor – to fight pathogenic microbes, mitigating infection and allowing damaged tissue to heal.<sup>9 10 11</sup>

Physically, beta-glucan creates a gel, slowing the transit of feed through the gut and allowing starches to be digested earlier in the system, thereby reducing the negative effects of starch in the hind gut.<sup>12 13</sup>

Freedom Health uses beta-glucan derived from oats,  $\beta(1,3/1,4)$ , as well as yeast  $\beta(1,3)$ . The beta-glucan from oats is mainly soluble, while the beta-glucan from yeast is mainly insoluble. Beta glucans of the form  $\beta(1,3)$  have the greatest immunomodulatory effect.

## Glutamine (outside E.U.)

Glutamine is the most abundant amino acid in humans as well as horses. Glutamine deficits can result in diarrhea, villous atrophy, mucosal ulceration, increased intestinal permeability and necrosis.<sup>14</sup>

Glutamine is a muscle fuel and also supplies nitrogen to the immune cells of the intestinal mucosa, which help to prevent pathogenic organisms from entering the circulatory system. Glutamine is considered to be a nonessential amino acid under normal conditions, because the body can create as much as is needed from scratch. But when the digestive system is stressed – for instance by ulcers or malnutrition – large amounts of glutamine are consumed, and supplements may be needed to replenish the supply.<sup>15 16</sup>

## Arginine (E.U. Only)

Arginine is a conditionally essential amino acid, associated with a healthy gut lining and rapid wound healing. Arginine is a precursor to nitric oxide (NO), which dilates blood vessels and improves circulation.<sup>17</sup> This improved circulation has been shown to enhance reproductive performance in mares and other animals, shortening the gestation period and increasing uterine arterial blood capacity.<sup>18</sup>

Arginine, along with glutamine, enhances gut function.<sup>19</sup> Arginine also decreases mucosal injury following pathogenic invasion and shortens recovery time following illness or intestinal wounding.<sup>20</sup> It stimulates the production of T cells, enhancing the immune response.<sup>21,22</sup>

Importantly from the viewpoint of performance athletes, arginine increases the production of growth hormones and insulin, leading to enhanced strength and stamina.<sup>23,24</sup>

## Threonine

Threonine is an essential amino acid and an important component of the mucin produced by the goblet cells distributed throughout the intestinal tract. By assisting metabolism and nutrient absorption, threonine contributes to a smoothly functioning GI tract. A deficiency of threonine slows the regeneration of the gut wall and depresses the production of mucus.<sup>25 26</sup>

Threonine is especially useful for wound healing and for treating stress, but it is also an essential link in the production of immunoglobulins.<sup>27</sup>

## Nucleotides

Although the GI tract is covered in mucus, the acids and enzymes nevertheless take a toll on the enterocytes lining the gut wall. The cells there are constantly dividing, ultimately managing to totally replace the intestinal lining about every three days.

This continuous cell division requires the replication of millions of DNA molecules every second. In turn, each DNA molecule is made up of several billion nucleotides. This represents an impressive amount of energy-intensive chemical synthesis. Clearly, maintaining an adequate level of nucleotides is a major, ongoing problem for the digestive system.

In general, DNA is synthesized through a complicated pathway that creates fresh nucleotides de novo. However, in the presence of pre-made nucleotides, the body can down-regulate this synthesis and instead use an enzyme named HGPRT to scavenge the intact nucleotides, improving the efficiency of cell repair.

In addition to simple maintenance, cell division is also critical for the repair of damaged tissue, including ulcers. DNA synthesis is thus a limiting factor in the healing process as well.

In times of stress, certain cells of the digestive system – including the mucus-producing goblet cells – are incapable of meeting the increased demand for nucleotides. Under these conditions, nucleotide supplements in the diet have been proven to be beneficial. Studies have shown that nucleotide supplements increase mucosal thickness and protein levels as well as speeding up intestinal recovery after chronic diarrhea and intestinal damage.<sup>28</sup>

Nucleotide supplements have been shown to increase the maturity and growth of normal enterocytes while reducing their dependence on exogenous glutamine.<sup>29</sup> The mechanism for this is not totally understood, but for intestinal villi to grow, stem cells in the crypts must divide and push their way up the length of the villi.<sup>30</sup> Exogenous DNA may enhance this process, perhaps explaining the extra crypt depth and increased surface area noted with nucleotide supplements.

Dietary nucleotides also seem to have an important beneficial effect on the intestinal microflora, stimulating the growth of beneficial bacteria and inhibiting pathogens. This may be due to yet another effect of dietary nucleotides reported in a NASA study and elsewhere: stimulation of the immune system.<sup>31 32</sup> In particular, lymphocytes and erythrocytes are not able to synthesize the purine-based nucleotides at all. For these cells, available nucleotides are essential to proper functioning.

## Yeast Products

*Saccharomyces cerevisiae* is a species of yeast with a long history in health and gastronomy. It has several beneficial properties, two of which are incorporated into SUCCEED DCP.

### **Yeast Beta-glucan**

As discussed earlier in the section on beta-glucan, yeast beta-glucans are of the form  $\beta(1,3)$  and are potent immunomodulators. Yeast beta glucan is extracted from the inner cell wall of the yeast cell. It is also an efficient mycotoxin adsorbent, helping to minimize the effect of mycotoxins produced from fungi and molds found in contaminated grain, forage and feeds.

### **Mannan Oligosaccharides**

Mannan oligosaccharide (MOS) is a polysaccharide composed of chains of the sugar mannose. It is extracted from the outer cell wall of *S. cerevisiae*. It binds to projections called fimbria on bacteria such as salmonella and clostridia.<sup>33</sup> Thus bound, fimbria can't form connections with enterocytes, preventing infection. The bound bacteria are then flushed from the system.<sup>34</sup> The result is a positive modulation of gut microflora and reduced risk of infection.

MOS can be digested by the enzymes of certain beneficial bacteria. So, in addition to discouraging pathogenic bacteria, MOS also promotes the growth of beneficial microbes such as lactobacillus.

MOS stimulates the immune system<sup>35</sup> and encourages the growth of intestinal villi, showing improved digestion and absorption of nutrients in various animal studies.<sup>36 37</sup> MOS also enhances the immune system by raising the levels of plasma and colostral IgG and bile IgA antibodies. Passive transfer of immunity within the first 24 hours of the foal's life is crucial for protection against pathogens, which may otherwise result in diarrhea, sepsis and even death.

Freedom Health uses MOS extracted from dried yeast.

## References

- 
- <sup>1</sup> Martin GP, Marriott C, Kellaway IW. *The interaction of progesterone with mucus glycoproteins*. Pharm Acta Helv. 1981;56(1):5-8.
  - <sup>2</sup> Kiviluoto T, Paimela H, Mustonen H, Kivilaakso E. *Exogenous surface-active phospholipid protects Necturus gastric mucosa against luminal acid and barrier-breaking agents*. Gastroenterology. 1991 Jan;100(1):38-46.
  - <sup>3</sup> McNeil PL, Ito S. *Gastrointestinal cell plasma membrane wounding and resealing in vivo*. Gastroenterology. 1989 May;96(5 Pt 1):1238-48.
  - <sup>4</sup> Kreitler B. *Feed and Nutrition: Fat: The Next Nutraceutical?* Thoroughbred Times, April 12, 2003.
  - <sup>5</sup> Davidson MH, Dugan LD, Burns JH, et al. *The hypocholesterolemic effects of beta-glucan in oatmeal and oat bran. A dose-controlled study*. JAMA 1991;265:1833-9.
  - <sup>6</sup> Bell S, Goldman VM, Bistran BR, et al. *Effect of beta-glucan from oats and yeast on serum lipids*. Crit Rev Food Sci Nutr 1999;39:189-202 [review].

- 
- <sup>7</sup> Braaten JT, Scott FW, Wood PJ, et al. *High beta-glucan oat bran and oat gum reduce postprandial blood glucose and insulin in subjects with and without type 2 diabetes.* Diabet Med 1994;11:312-8.
- <sup>8</sup> Tappy LE, Gugolz E, et al. *Effects of breakfast cereals containing various amounts of beta-glucan fibers on plasma glucose and insulin responses in NIDDM subjects.* Diabetes Care 19(8):831-4.
- <sup>9</sup> Czop JK. *The role of beta-glucan receptors on blood and tissue leukocytes in phagocytosis and metabolic activation.* Pathol Immunopathol Res 1985;5:286-96.
- <sup>10</sup> Estrada A, Yun CH, Van Kessel A, et al. *Immunomodulatory activities of oat beta-glucan in vitro and in vivo.* Microbiol Immunol 1997;41:991-8.
- <sup>11</sup> Reid DM, Montoya M, et al. *Expression of the beta-glucan receptor, Dectin-1, on murine leukocytes in situ correlates with its function in pathogen recognition and reveals potential roles in leukocyte interactions.* J Leukoc Biol 76(1):86-94.
- <sup>12</sup> Bohm N, Kulicke W. *Rheological studies of barley (1-3)(1-4) beta-glucan in concentrated solution,* Carbohydrate Research, 1999, 315, 293-301.
- <sup>13</sup> Wursch P, Sunyer FX. *The role of viscous soluble fiber in the metabolic control of diabetes. A review with special emphasis on cereals rich in beta-glucan.* Diabetes Care 20(11):1774-80.
- <sup>14</sup> Roth E, Spittler A, Oehler R. *Glutamine: effects on the immune system, protein balance and intestinal functions.* Wien Klin Wochenshr. 1996;108(21):667-8.
- <sup>15</sup> Duckworth DH, Madison JB, et al. *Arteriovenous differences for glutamine in the equine gastrointestinal tract.* Am J Vet Res 53(10): 1864-7.
- <sup>16</sup> Vazquez P, Gomez de Segura IA, Cos A, Candela CG, De Miguel E. *Response of the intestinal mucosa to different enteral diets in situations of surgical stress and malnutrition.* Nutr Hosp. 1996 Nov-Dec;11(6):321-7.
- <sup>17</sup> Moncada, Salvador, and Annie Higgs. "The L-Arginine-Nitric Oxide Pathway." New England Journal of Medicine 329, no. 27 (December 30, 1993): 2002–12. doi:10.1056/NEJM199312303292706.
- <sup>18</sup> Mortensen, Chris J., Dale E. Kelley, and Lori K. Warren. "Supplemental L-Arginine Shortens Gestation Length and Increases Mare Uterine Blood Flow before and after Parturition." *Journal of Equine Veterinary Science* 31, no. 9 (September 1, 2011): 514–20. doi:10.1016/j.jevs.2011.01.004.
- <sup>19</sup> Stechmiller, J. K., D. Treloar, and N. Allen. "Gut Dysfunction in Critically Ill Patients: A Review of the Literature." *American Journal of Critical Care* 6, no. 3 (May 1, 1997): 204–9.
- <sup>20</sup> Sukhotnik, Igor, Jorge Mogilner, Michael M. Krausz, Michael Lurie, Mark Hirsh, Arnold G. Coran, and Eitan Shiloni. "Oral Arginine Reduces Gut Mucosal Injury Caused by Lipopolysaccharide Endotoxemia in Rat." *Journal of Surgical Research* 122, no. 2 (December 1, 2004): 256–62. doi:10.1016/j.jss.2004.07.004.

- 
- <sup>21</sup> Madden, Hugh P., Robert J. Breslin, Hannah L. Wasserkrug, Gershon Efron, and Adrian Barbul. "Stimulation of T Cell Immunity by Arginine Enhances Survival in Peritonitis." *Journal of Surgical Research* 44, no. 6 (June 1, 1988): 658–63. doi:10.1016/0022-4804(88)90097-2.
- <sup>22</sup> Gianotti, L, J W Alexander, T Pyles, and R Fukushima. "Arginine-Supplemented Diets Improve Survival in Gut-Derived Sepsis and Peritonitis by Modulating Bacterial Clearance. The Role of Nitric Oxide." *Annals of Surgery* 217, no. 6 (June 1993): 644–54.
- <sup>23</sup> Sureda, Antoni, and Antoni Pons. "Arginine and Citrulline Supplementation in Sports and Exercise: Ergogenic Nutrients?" 59 (2012): 18–28. doi:10.1159/000341937.
- <sup>24</sup> Schmidt, Harald H. H. W., Timothy D. Warner, Kunio Ishii, Hong Sheng, and Ferid Murad. "Insulin Secretion from Pancreatic B Cells Caused by L-Arginine-Derived Nitrogen Oxides." *Science* 255, no. 5045 (1992): 721–23.
- <sup>25</sup> Bertolo RF, Chen CZ, Law G, Pencharz PB, Ball RO. *Threonine requirement of neonatal piglets receiving total parenteral nutrition is considerably lower than that of piglets receiving an identical diet intragastrically.* J Nutr. 1998 Oct;128(10):1752-9.
- <sup>26</sup> Ball RO, Law G, Bertolo RFP, Pencharz PB. *Adequate oral threonine is critical for mucin production and mucosal growth by neonatal piglet gut.* Proceedings of the VIIIth International Symposium on Protein Metabolism and Nutrition, EAAP, 1999
- <sup>27</sup> Cuaron JA, Chapple RP, Easter RA. *Effect of lysine and threonine supplementation of sorghum in gestation diets on nitrogen balance and plasma constituents in first litter gilts.* J. Anim. Sci., 58, 631-637
- <sup>28</sup> Bueno J, Torres A, Almendros A, Carmona R, Nunez MC and Gil A, (1994) *Effect of dietary nucleotides on small intestinal repair after diarrhea. Histological and ultrastructural changes.* Gut 35: 926-933.
- <sup>29</sup> Uauy R, Stringel G, Thomas R and Quan R, (1990) *Effect of dietary nucleosides on growth and maturation of the developing gut in the rat.* J. Pediatr. Gastroenteral. Nutr. 10:497-503.
- <sup>30</sup> Marshman E., Booth C., Potten CS., *The intestinal epithelial stem cell.* Bioessays 2002 Jan; 24(1):91-8.
- <sup>31</sup> Hales, N. *Diet Supplement May Help Boost Immune System.* American Society for Parenteral and Enteral Nutrition 25<sup>th</sup> Clinical Congress in Chicago.
- <sup>32</sup> Lin, Cheng-mao. *Effect of Dietary Nucleotide Supplementation on In Vivo and In Vitro Immune Function in Protein-Malnourished Mice.* University of Florida. PhD. Dissertation. December 1995.
- <sup>33</sup> Ip WK, Lau YL. Role of mannose-binding lectin in the innate defense against *Candida albicans*: enhancement of complement activation, but lack of opsonic function, in phagocytosis by human dendritic cells. *J Infect Dis* 2004 Aug 1;190(3):632-40. Epub 2004 Jun 28.
- <sup>34</sup> Swanson KS, Grieshop CM, Flickinger EA, Healy HP, Dawson KA, Merchen NR, Fahey GC Jr. *Effects of supplemental fructooligosaccharides plus mannanoligosaccharides on*

---

*immune function and ileal and fecal microbial populations in adult dogs.* Arch Tierernahr. 2002 Aug;56(4):309-18.

<sup>35</sup> Bland EJ, Keshavarz T, Bucke C. *The influence of small oligosaccharides on the immune system.* Carbohydrate Research, vol 339, issue 10.

<sup>36</sup> Newman, K. 1994. *Mannan-oligosaccharides: Natural polymers with significant impact on the gastrointestinal microflora and the immune system.* Biotechnology in the Feed Industry, Nottingham University Press, Nottingham, UK, pp. 167-174.

<sup>37</sup> Davis E, Maxwell C, Kegley B, de Rodas B, Friesen K and Hellwig D, *Efficacy of Mannan Oligosaccharide (Bio-Mos) Addition at Two Levels of Supplemental Copper on Performance and Immunocompetence of Early Weaned Pigs.* Arkansas Animal Science Department Report 1999.